

FILE COPY

2

SECURITY

AD-A201 768 IDENTIFICATION PAGE

Form Approved
OMB No 0704-0188
Exp Date Jun 30, 1986

1a. RE L		1b. RESTRICTIVE MARKINGS	
2a. SECURITY CLASSIFICATION AUTHORITY UNCLASSIFIED		3. DISTRIBUTION / AVAILABILITY OF REPORT	
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE UNCLASSIFIED			
4. PERFORMING ORGANIZATION REPORT NUMBER(S)		5. MONITORING ORGANIZATION REPORT NUMBER(S)	
6a. NAME OF PERFORMING ORGANIZATION DIV OF ET & BICM	6b. OFFICE SYMBOL (If applicable)	7a. NAME OF MONITORING ORGANIZATION WRAIR	
6c. ADDRESS (City, State, and ZIP Code) Washington, DC 20307-5100		7b. ADDRESS (City, State, and ZIP Code) Washington, DC 20307-5100	
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Ft Detrick, Frederick, MD	8b. OFFICE SYMBOL (If applicable)	9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) US Army Medical Res & Dev Command Ft Detrick, Frederick, MD 21701-5012		10. SOURCE OF FUNDING NUMBERS	
		PROGRAM ELEMENT NO.	PROJECT NO.
		TASK NO.	WORK UNIT ACCESSION NO.
11. TITLE (Include Security Classification) LEISHMANIA MEXICANA: UPTAKE OF SODIUM STIBOGLUCONATE (PENTOSTAM) AND PENTAMIDINE BY PARASITE AND MACROPHAGES			
12. PERSONAL AUTHOR(S) J.D. BERMAN, J.V. GALLALEE, B.D. HANSEN			
13a. TYPE OF REPORT Manuscript	13b. TIME COVERED FROM TO	14. DATE OF REPORT (Year, Month, Day)	15. PAGE COUNT
16. SUPPLEMENTARY NOTATION			
17. COSATI CODES		18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)	
FIELD	GROUP SUB-GROUP		
19. ABSTRACT (Continue on reverse if necessary and identify by block number)			
<div style="text-align: right;"> <p>DTIC ELECTE OCT 11 1988 D</p> </div>			
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS		21. ABSTRACT SECURITY CLASSIFICATION	
22a. NAME OF RESPONSIBLE INDIVIDUAL JONATHAN D. BERMAN		22b. TELEPHONE (Include Area Code)	22c. OFFICE SYMBOL

DD FORM 1473, 84 MAR

83 APR edition may be used until exhausted

All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE
UNCLASSIFIED

88 10 6 05 9

RESEARCH BRIEF

Leishmania mexicana: Uptake of Sodium Stibogluconate (Pentostam) and Pentamidine by Parasite and Macrophages

JONATHAN D. BERMAN,* JAMES V. GALLALEE,* AND BRIAN D. HANSEN†

Divisions of *Experimental Therapeutics and †Biochemistry, Walter Reed Army Institute of Research, Walter Reed Army Medical Center, Washington, DC 20307-5100, U.S.A.

(Accepted for publication 16 December 1986)

BERMAN, J. D., GALLALEE, J. V., AND HANSEN, B. D. 1987. *Leishmania mexicana*: Uptake of sodium stibogluconate (Pentostam) and pentamidine by parasite and macrophages. *Experimental Parasitology* 64, 127-131. © 1987 Academic Press, Inc.

Pentavalent antimonials in the form of sodium stibogluconate (Pentostam) or meglumine antimonate (Glucantime) are the primary therapeutic agents for leishmaniasis. Pentamidine is clinically effective but has been relegated to the status of a secondary agent because of toxicity. However, pentamidine is a widely employed antimicrobial because it is also efficacious in *Pneumocystis carinii* infections and in African trypanosomiasis. There are two stages of *Leishmania* spp.: the mammalian form (amastigote) which is an obligate intramacrophage microorganism and the insect vector form (promastigote). Both Pentostam and pentamidine are more active against intramacrophage amastigotes than extracellular promastigotes (Berman *et al.* 1980), and the fact that both drugs are clinical agents indicates that in general both drugs are more active against amastigotes than mammalian cells.†

Uptake of drugs by cells can be via simple diffusion, facilitated diffusion, or active transport. Acquisition of radiolabeled Pentostam and pentamidine permitted this report of the uptake of these drugs by *Leishmania* spp. amastigotes and *Leishmania* spp. promastigotes and macrophages.

The strain used in these experiments, *L. mexicana* WR 227, was cultured from a patient who acquired cutaneous infection in Panama. The strain was determined to be *L. mexicana* by isoenzyme analysis (R. D. Kreutzer, personal communication). Promastigotes of WR 227 were maintained in Schneider's drosophila medium revised (GIBCO, Grand Island, NY, U.S.A.) with the addition of 20% heat inactivated fetal calf serum (Hendricks and Wright 1979). Promastigotes were grown to late log phase ($40 \times 10^6/\text{ml}$), washed with HBSS (GIBCO), and resuspended in HBSS for experimentation. Amastigotes of WR 227 were obtained from amastigote-containing J774 mac-

rophages by the method of Chang (1980) as modified (Berman *et al.* 1985). The macrophages were infected with promastigotes, maintained until the promastigotes transformed to amastigotes, and lysed to release the intracellular amastigotes. The amastigotes, purified by Percoll sedimentation, were washed and resuspended in HBSS for experimentation. J774 macrophages were also washed and resuspended in HBSS for experimentation. Experiments with all cell types were performed at the pH of HBSS (pH 7.4) because this pH is optimal for the macrophages and adequate for promastigote viability. The pH of the phagolysosome in which amastigotes reside is not known.

In experiments involving uptake of drug versus time, cells in HBSS were incubated with 1.6 mM ^{125}Sb in the form of Pentostam (^{125}Sb labeled) or 6.7 μM [^3H]pentamidine isethionate ([^3H]pentamidine). At appropriate time points, 0.2 ml of cell suspension containing 10×10^6 amastigotes, 14×10^6 promastigotes, or 3×10^5 macrophages was withdrawn and centrifuged ($12,000g$ for 3 min) through silicone F-50 oil (General Electric, Waterford, NY, U.S.A.), and the pellet was scintillation counted. In experiments involving velocity of drug uptake versus drug concentration, the same number of cells in 0.2 ml was exposed to 1 mM ^{125}Sb and 0-49 mM cold Sb in the form of Pentostam or to 3 μM [^3H]pentamidine and 0-147 μM cold pentamidine, for 1 min. The cells were then pelleted and counted. Experiments with amastigotes and macrophages were performed at 35 C; experiments with promastigotes were performed at 26 C. Cellular volumes were approximated by comparing the volume of a large number of pelleted cells to known volumes of water. Protein determinations were made by the Lowry technique. The amount of 1×10^6 amastigotes constitutes 0.027 μl and 3.1 μg protein; 1

TABLE I
In Vitro Antileishmanial Activity of [125 Sb]Pentostam and Unlabeled Pentostam

Pentostam	Concn (mM)	Promastigotes: % control 2 days after drug exposure	Amastigotes: % control 2 days after drug exposure
Cold Sb	1.2	90 ^a	75 ^a
	2.4	74 ^a	62 ^a
	4.0	60 ^a	39 ^a
125 Sb	1.2	76–85 ^b	72–75 ^b
	2.4	64–71 ^b	62–63 ^b
	4.0	42–55 ^b	39–48 ^b

Note. *Leishmania mexicana* WR 227 promastigotes or amastigotes were exposed to cold or radiolabeled Sb in the form of Pentostam for 4 hr, washed, and resuspended in promastigote culture media (Berman *et al.* 1985). After 2 days, the number of promastigotes in each drug treated culture was expressed as a percentage of the number of organisms in simultaneously cultivated controls.

^a Values from Berman *et al.* 1985.

^b Range of two experiments.

$\times 10^6$ promastigotes constitute 0.10 μ l and 7.0 μ g protein; 1×10^6 J774 macrophages constitute 20 μ l and 300 μ g protein.

[125 Sb]Sodium stibogluconate was custom synthesized by Amersham Corp. (Chicago, IL, U.S.A.) under the direction of Burroughs Wellcome Co. (London, UK) and had a specific activity of 0.375 mCi/mmol Sb. Because cold Pentostam consists of an unknown number of uncharacterized combinations of Sb and carbohydrates derived from gluconic acid, the purity of the radiolabeled product could not be compared to that of the cold drug by standard chemical means. Instead, the purity of the radiolabeled product was determined by bioassay using previously published procedures (Berman *et al.* 1985). The anti-leishmanial activity of radiolabeled Pentostam was similar to that of cold Pentostam for both amastigotes and promastigotes (Table I).

[3 H]Pentamidine isethionate (310 mCi/mmol) was custom synthesized by Moravsek Biochemicals, La-Brea, CA, U.S.A. and its purity was verified by HPLC (C. Dickenson, personal communication). Quantities of pentamidine isethionate refer to quantities of the total salt; 57% of pentamidine isethionate is pentamidine base. Cold Pentostam was obtained from Burroughs Wellcome. Cold pentamidine isethionate was obtained from May & Baker (Dagenham, UK).

The uptake of 1.6 mM [125 Sb]Pentostam as a function of time by *L. mexicana* amastigotes, *Leishmania mexicana* promastigotes, and J774 macrophages is seen in Fig. 1a. During the first minute of drug exposure, there was an initial rapid uptake by three cell types. By 3 min of exposure, amastigotes concentrated Sb more than macrophages. By the end of the hour of incubation, the Sb concentration within amastigotes (49 mM: Concentration factor ≈ 30 compared

to the external medium) was about three times that of both promastigotes and macrophages (14 mM: Concentration factor ≈ 9). Croft *et al.* (1981) found that after 24 hr, amastigotes and promastigotes concentrated [125 Sb]Pentostam by factors of 10–20 and 2–10, respectively. The uptake of [125 Sb] as a function of concentration is seen in Fig. 1b. Although uptake at less than 1 min was not ascertained, the generally linear increase in uptake by all three cell types in these 1-min experiments ($R^2 \geq 0.97$ for each straight line) indicates that Pentostam is taken predominately up by cells by simple diffusion. Therefore, the ability of all cell types tested to concentrate Pentostam may be due to binding of the drug to macromolecules in the cells.

The uptake of 6.7 μ M [3 H]pentamidine as a function of time is seen in Fig. 2a. As with Pentostam, the [3 H]pentamidine uptake was initially greater for amastigotes and macrophages than for promastigotes (Fig. 2a). By the end of the hour of incubation, pentamidine was concentrated within amastigotes by a factor of 20 compared to the external medium and within macrophages and promastigotes by factors of 6.3 and 4.5 compared to the external medium. The uptake of pentamidine versus concentration indicates that drug uptake is predominantly by simple diffusion ($R^2 \geq 0.98$ for the straight lines in Fig. 2b). Thus, concentration of this drug with cells is probably due also to binding to macromolecules. The data with pentamidine are consistent with the previous determination of pentamidine uptake by *Trypanosoma brucei* stains. In one report, pentamidine was concentrated by a factor of 2.3–17 (Damper and Patton 1976a); in another report, the drug was taken up by linear diffusion by some of the strains tested (Damper and Patton 1976b).

The greater ability of amastigotes, compared to promastigotes and mammalian macrophages, to concentrate Pentostam and the generally greater ability of




FIG. 1. Uptake of radiolabeled drugs by *Leishmania mexicana* amastigotes (AMAS), *L. mexicana* promastigotes (PROS), and J774 macrophages (MAC) over 60 min. (a) Uptake of $[^{125}\text{Sb}]$ Pentostam. (b) Uptake of $[^3\text{H}]$ pentamidine.




FIG. 2. Uptake of radiolabeled drugs by *Leishmania mexicana* amastigotes (AMAS) and promastigotes (PROS) and by J774 macrophages (MAC), versus concentration. (a) Uptake of $[^{125}\text{Sb}]$ Pentostam. (b) Uptake of $[^3\text{H}]$ pentamidine.

amastigotes to concentrate pentamidine suggest a mechanism that may contribute to the toxicity of these drugs to amastigotes compared to promastigotes and mammalian cells.

(We thank J. Best for his expert technical assistance.)

REFERENCES

- BERMAN, J. D., DWYER, D. W., AND WYLER, D. J. 1980. An *in vitro* model for investigation of chemotherapeutic agents in leishmaniasis. *Journal of Infectious Diseases* 142, 83-86.
- BERMAN, J. D., WADDELL, D., AND HANSON, B. D. 1985. Biochemical mechanism of the antileishmanial activity of sodium stibogluconate. *Antimicrobial Agents Chemotherapy* 27, 916-920.
- CHANG, K.-P. 1980. Human cutaneous *Leishmania* in a mouse macrophage line: Propagation and isolation of intracellular parasites. *Science* 209, 1240-1242.
- CROFT, S. L., NEAME, K. D., AND HOMEWOOD, J. 1980. Human cutaneous *Leishmania* in a mouse macrophage line: Propagation and isolation of intracellular parasites. *Science* 209, 1240-1242.

- C. A. 1981. Accumulation of [125 Sb]sodium stibogluconate by *Leishmania mexicana mazonensis* and *Leishmania donovani* in vitro. *Comparative Biochemistry and Physiology* 68c, 95-98.
- DAMPER, D., AND PATTON, C. L. 1976a. Pentamidine transport in *Trypanosoma brucei*—Kinetics and specificity. *Biochemical Pharmacology* 25, 271-276.
- DAMPER, D., AND PATTON, C. L. 1976b. Pentamidine transport and sensitivity in brucei-group *Trypanosoma*. *Journal of Protozoology* 23, 349-356.
- HENDRICKS, L. D., AND WRIGHT, N. 1979. Diagnosis of cutaneous leishmaniasis by *in vitro* cultivation of saline aspirates in Schneider's drosophila medium. *American Journal of Tropical Medicine and Hygiene* 28, 962-964.

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	20

